

2002 DEC 27 AM 11: 24

# HIGH PRODUCTION VOLUME (HPV) CHALLENGE PROGRAM HYDROQUINONE MONOMETHYL ETHER CATEGORY TEST PLAN

4-HYDROXYANISOLE (CAS No. 150-76-5)

PREPARED BY:

MEMBER COMPANIES OF THE AMERICAN CHEMISTRY COUNCIL'S HYDROQUINONE PRECURSORS AND DERIVATIVES PANEL HYDROQUINONE MONOMETHYL ETHER TASK FORCE

EASTMAN CHEMICAL COMPANY
RHODIA INCORPORATED
SPECIALTYCHEM PRODUCTS CORPORATION
BORREGAARD

December 20, 2002

## 4-HYDROXYANISOLE BACKGROUND INFORMATION

### 4-Hydroxyanisole CAS No.: 150-76-5 Background Information, Test Plan, and Robust Summaries

#### Introduction

This test plan is provided by the American Chemistry Council HQMME Task Force of the HQPD Panel, in fulfillment of its commitment under the US HPV Chemical Challenge Program. Members of the HQMME Task Force are Rhodia, Inc., Eastman Chemical Company, Borregaard, and Specialtychem Products Corporation.

#### Summary

4-Hydroxyanisole has been manufactured in the US and abroad for many years. It is used primarily in industrial settings as an oxygen scavenger for polymerization inhibition in the manufacture of a broad array of products. Non-industrial applications for this material are very minor. A small quantity of 4-Hydroxyanisole is manufactured by organizations – not part of the HQMME Task Force – for pharmaceutical use. As shown in the Test Plan, the Physical/chemical Properties, Environmental Fate and Effects, and Mammalian Toxicity associated with 4-hydroxyanisole are well characterized. No further testing has been proposed for this material. Robust summaries are provided as Appendix A.

#### Chemical Identity:

4-Hydroxyanisole CAS No.: 150-76-5

#### **Background Information: Manufacturing and Commercial Applications**

#### <u>Manufacturing</u>

4-Hydroxyanisole is made in batch processes using hydroquinone, sodium hydroxide and methyl chloride as the primary starting materials. This reaction produces crude 4-hydroxyanisole and a minor co-product 1,4-dimethoxybenzene. The crude 4-hydroxyanisole is distilled to remove trace impurities.

#### **Commercial Applications**

4-Hydroxyanisole has been manufactured in the United States for nearly 40 years, and is widely used throughout the chemical industry due to its excellent stability and functionality as

an oxygen scavenger. The majority of all usage is in industrial applications as a polymerization inhibitor for acrylic acid and esters, acrylonitrile and methacrylic acid and esters. It is also utilized as the starting material for the production of butylated hydroxyanisole. A very minor application that has only just recently been developed is as an active ingredient in a dermatological preparation used to lighten skin. The safety of the product for this consumer use was reviewed by the FDA and included data were not completely accessible to the Panel.

#### Shipping/Distribution

4-Hydroxyanisole is shipped extensively throughout the world from manufacturing plants located in the United States, Western Europe, Japan and India.

#### **Worker Exposure**

The synthetic fine organic batch manufacturing industry has a long safety record and only sophisticated industrial users handle this material. Exposure of workers to 4-hydroxyanisole is likely highest in the area of material solidification and packaging rather than from chemical manufacturing. This material is made as a molten material that is usually flaked or cast and then re-melted or dissolved at the user's facility. As there is little dust generation with the use of this material, the main exposure is from vapor generated in the hot liquid operations as it is solidified and packaged. This is easily controlled by any number of air control technology applications and good industrial hygiene practices by workers. There is a TWA exposure level of 5 mg/m³ set by the ACGIH and by NIOSH.

#### Consumer exposure

Consumer exposure is minimal except through its use as an active agent in a dermatological product recently approved by the US FDA. This product is listed by the FDA as being available for use by prescription only. The product is applied directly to the skin in a cream/ointment form.

#### **HQMME Test Plan**

As shown in the Test Plan below, the Physical/chemical properties, Environmental Fate and Effects, and Mammalian Toxicity associated with 4-hydroxyanisole are well characterized. No further testing has been proposed for this material.

			Physi	cal-Ch	nemica	ıl		
Melting Point	Boiling	Boiling Point		Vapor Pressure		Partitio Coefficie		Water Solubility
А	A	Α		NA		Α		А
			Enviro	nmen	tal Fat	:e		
Photodegradation		Stability in Water		Transport/ Distribution			Biodegradation	
Calc		١	IA AI	Calc		Α		
			Ed	otoxi	city			
Acute Toxicity to Fish			Acute Toxicity to Aquatic Plants (e.g., Algae)				Acute Toxicity to Aquatic Invertebrates (e.g., Daphnia)	
А			A				A	
			Mamm	alian <sup>*</sup>	Toxicit	:у		
Acute Toxicity	Bacterial Genetic Toxicity <i>In</i> <i>Vitr</i> o		Mammaliar Genetic Toxicity <i>In</i> <i>Vitr</i> o	Genetic D oxicity <i>In</i> To		Reproductive Toxicity		Developmental Toxicity
А	А		Α		Α	А		А

Legend					
Symbol	Description				
Calc	Endpoint requirement fulfilled based on calculated data				
Α	Endpoint requirement fulfilled with adequate existing data				
NR	Not required per the OECD SIDS guidance				
NA	Not applicable due to physical/chemical properties				